

How to calculate spectra of Rabi and related models?

Andrzej J. Maciejewski*

J. Kepler Institute of Astronomy, University of Zielona Góra, Licealna 9, PL-65-417 Zielona Góra, Poland.

Maria Przybylska†

Institute of Physics, University of Zielona Góra, Licealna 9, 65-417 Zielona Góra, Poland

Tomasz Stachowiak‡

Center for Theoretical Physics PAS, Al. Lotników 32/46, 02-668 Warsaw, Poland

(Dated: November 7, 2012)

We show how to use properly the Bargmann space of entire functions in the analysis of the Rabi model. We are able to correct a serious error in recent papers on the topic and develop a corrected method of finding the spectrum applicable also to other, more general systems.

PACS numbers: 03.65.Ge, 02.30.Ik, 42.50.Pq

Keywords: Rabi model

Several papers devoted to determination of the spectrum of the Rabi model have appeared recently, see e.g. [1–3], and references therein for a description of various approaches to this problem. Let us recall that the Rabi model describes interaction of a two-level atom with a single harmonic mode of the electromagnetic field. In the Bargmann-Fock representation this model is described by the following system of two differential equations

$$\begin{aligned}(z + \lambda) \frac{d\psi_1}{dz} &= (E - \lambda z) \psi_1 - \mu \psi_2, \\ (z - \lambda) \frac{d\psi_2}{dz} &= (E + \lambda z) \psi_2 - \mu \psi_1,\end{aligned}\quad (1)$$

where E is the energy, λ is the atom-field coupling constant, and 2μ is the atomic level separation. In this representation, two component wave function $\psi = (\psi_1, \psi_2)$ is an element of Hilbert space $\mathcal{H}^2 = \mathcal{H} \times \mathcal{H}$, where $\mathcal{H}$ is the Bargmann-Fock Hilbert space of entire functions of one variable $z \in \mathbb{C}$. The scalar product in $\mathcal{H}$ is given by

$$\langle f, g \rangle = \frac{1}{\pi} \int_{\mathbb{C}} \overline{f(z)} g(z) e^{-|z|^2} d(\Re(z)) d(\Im(z)).$$

An entire function $f(z)$ belongs to $\mathcal{H}$ if it has the proper growth at infinity, for details, see [4]. Thus, energy E belongs to the spectrum of the problem, if and only if for this value of E equations (1) have entire solution $\psi = (\psi_1, \psi_2)$ with the proper behaviour at infinity. However, for the considered system all entire solutions belong to $\mathcal{H}^2$, see [5] for an explanation.

Equations (1) have two singular regular points at $z = \lambda$ and $z = -\lambda$, while infinity is an irregular singular point. The system has a $\mathbb{Z}_2$ symmetry. It is invariant with respect to the involution $\tau : \mathcal{H}^2 \rightarrow \mathcal{H}^2$ given by $\tau(\psi_1, \psi_2)(z) = (\psi_2(-z), \psi_1(-z))$. In other words, if $(\psi_1(z), \psi_2(z))$ is a solution of this system, then also $(\psi_2(-z), \psi_1(-z))$ is its solution. We say that a solution $\psi = (\psi_1, \psi_2)$ of (1) has parity $\sigma \in \{-1, +1\}$, if

$$\tau(\psi) = \sigma \psi.$$

In [1] it was pointed out that the above mentioned symmetry plays the crucial role in determination of its spectrum. Later, in [3], it was shown that one can determine the spectrum of the Rabi model without any references to its $\mathbb{Z}_2$ symmetry. However, the method used in [3] has limited application as it is based on a result valid only for third order recurrence relations, see [6].

Braak's approach is very interesting because it gives a chance for generalisation beyond third order recurrence relations. However in its original formulation it contains a flaw which invalidates its results concerning the spectrum of the standard Rabi model. The aim of our paper is to correct results of [1] and to present our direct method which can be used to study of systems more complicated than the Rabi model.

We applied our method to study a generalised Rabi model with broken $\mathbb{Z}_2$ symmetry introduced in [1]. In the Bargmann representation it is described by the following equations

$$\begin{aligned}(z + \lambda) \frac{d\psi_1}{dz} &= (E - \epsilon - \lambda z) \psi_1 - \mu \psi_2, \\ (z - \lambda) \frac{d\psi_2}{dz} &= (E + \epsilon + \lambda z) \psi_2 - \mu \psi_1.\end{aligned}\quad (2)$$

For $\epsilon = 0$ it coincides with the Rabi model (1). We denote by $p := (E, \lambda, \mu, \epsilon)$ parameters of this model. Instead of E we will use also the parameter $x := E + \lambda^2$. Let us assume that x is not a non-negative integer. Then the spectrum of the problem coincides with zeros of one function $w(p)$ defined in the following way. Let

$$w(p; y) := H_1(y) H_2'(y) - H_1'(y) H_2(y), \quad (3)$$

where

$$H_1(y) := \text{HeunC}(a_0; y), \quad (4)$$

and

$$H_2(y) := \text{HeunC}(a_1; 1 - y), \quad (5)$$

are the confluent Heun functions, see [7], with parameters $a_0 := (\alpha, \beta, \gamma, \delta, \eta)$, and $a_1 := (-\alpha, \gamma, \beta, -\delta, \delta + \eta)$; in terms of $(x, \lambda, \mu, \epsilon)$ these parameters are given by $\alpha = 4\lambda^2$, $\beta = -x + \epsilon$, and

$$\begin{aligned} \gamma &= -1 - x - \epsilon, & \delta &= 2(1 - 2\epsilon)\lambda^2, \\ 2\eta &= 1 - 2\mu^2 + (1 + x)(x - 4\lambda^2) + \epsilon(1 + 4\lambda^2) - \epsilon^2. \end{aligned} \quad (6)$$

Then, $W(p) := w(p; 1/2)$.

Here we just recall the basic steps of the procedure which was formulated for the Rabi model (1). It is fully described in [8].

Let us introduce new variables $y := z + \lambda$, $\varphi_i(y) = e^{-\lambda y} \psi_i(y - \lambda)$, for $i=1,2$. It is obvious that functions $\psi_i(z)$ are entire if and only if functions $\varphi_i(y)$ are entire. In the new variables system (1) reads

$$\begin{aligned} y \frac{d\varphi_1}{dy} &= x\varphi_1 - \mu\varphi_2, \\ (y - 2\lambda) \frac{d\varphi_2}{dy} &= (x - 4\lambda^2 + 2\lambda y)\varphi_2 - \mu\varphi_1, \end{aligned} \quad (7)$$

Let us assume that for a given $p = (x, \lambda, \mu)$, system (7) has an entire solution $\varphi(y) = (\varphi_1(y), \varphi_2(y))$, where

$$\varphi_1(y) = \sum_{n=0}^{\infty} \tilde{\Phi}_n(p) y^n, \quad \varphi_2(y) = \sum_{n=0}^{\infty} \Phi_n(p) y^n. \quad (8)$$

To proceed further we assume also that $x \notin \mathbb{Z}$, since the existence of solutions for integer x has long been established, see [9]. Then we insert the expression for $\varphi_2(y)$ into the first equation (7), and integrate it. We obtain

$$\tilde{\Phi}_n(p) = \frac{\mu}{x - n} \Phi_n(p), \quad \text{for } n \in \mathbb{N}. \quad (9)$$

Function $\Phi_n(y)$ can be determined recursively. In this way we obtain a solution $\varphi(y)$, which is locally holomorphic at $y = 0$. Thus, we have a solution $\psi(z)$ which is locally holomorphic at $z = -\lambda$. However, we are looking for an entire solution. In order to find conditions which guarantee that $\psi(z)$ is entire we can assume, without loss of generality that it has a given parity σ . Thus putting $\psi_\sigma(z) = (\psi_{1,\sigma}(z), \psi_{2,\sigma}(z))$ we have $\psi_{1,\sigma}(z) = \sigma\psi_{2,\sigma}(-z)$ for all $z \in \mathbb{C}$. Hence, function

$$G_\sigma(z, p) := \psi_{2,\sigma}(-z) - \sigma\psi_{1,\sigma}(z), \quad (10)$$

must vanish identically for all $z \in \mathbb{C}$. In [8], Braak claims that

1. if $G_\sigma(z, p) = 0$ for a fixed but *arbitrary* z belonging to the intersection of domains of definiteness of $\psi_{1,\sigma}(z)$ and $\psi_{2,\sigma}(-z)$, then it vanishes identically.

That is, if for a given p there exists $z_\star \in D(-\lambda, 2|\lambda|) \cap D(\lambda, 2|\lambda|)$ such that $G_\sigma(z_\star, p) = 0$, then $G_\sigma(z, p) = 0$ for an arbitrary $z \in \mathbb{C}$. Here $D(z_0, r) \subset \mathbb{C}$ is an open disc of radius r and the centre at z_0 .

2. x belongs to the spectrum of the problem if and only if there exists $z_\star \in D(-\lambda, 2|\lambda|) \cap D(\lambda, 2|\lambda|)$ such that $G_\sigma(z_\star, p) = 0$, for a certain $p = (x, \lambda, \mu)$.

Both of these statements are incorrect. There exist entire functions, e.g., $\sin(z)$, which have infinite number of zeros but they are not zero functions. So, the first claim needs at least some justification. We show that in the considered context it is not valid. To see this it is enough to make a contour graph of function $G_\sigma(z, p)$ on the plane (x, z) , see Fig. 1. One can notice that there




FIG. 1. The zero level contours of $G_+(z, p)$ (continuous lines), and $G_-(z, p)$ (dashed lines) on the (x, z) plane, for $\lambda = 0.7$ and $\mu = 0.4$.

exist x such that $G_\sigma(z, p) = 0$ identically in z , as well as that there exist isolated zeros of $G_\sigma(z, p)$. An example is presented in Fig. 1. This also shows that the second claim of Braak is not correct.

We notice that the function $g(z) := G_\sigma(z, p)$, as a function of z , satisfies a second order linear differential equation. In fact, it can be checked directly that

$$(z^2 - \lambda^2)g'' + [z(1 - 2x - 3\lambda^2) - \lambda]g' + [\lambda z(1 - \lambda z) + (x - \lambda^2)^2 - \lambda^2 - \mu^2]g = 0. \quad (11)$$

So, fixing values $G_\sigma(z, p)$ and $G'_\sigma(z, p)$ at a given z , we determine $G_\sigma(z, p)$ uniquely. Hence, the spectrum of the Rabi problem in Braak's settings is determined by *two* transcendental equation $G_\sigma(z_\star, p) = 0$ and $G'_\sigma(z_\star, p) = 0$, not just one $G_\sigma(z_\star, p) = 0$, for a certain $z_\star$, as is claimed

in [1]. This is illustrated in Fig. 2. Here we just remark that choice $z_* = 0$ made in [1] was lucky because, for $x < 30$, all zeros of $G_\sigma(0, p)$ belong to the spectrum.




FIG. 2. Graphs of $G_+(z_*, p)$ (continuous line), and $G'_+(z_*, p)$ (dotted lines) for $z_* = -0.43$ as functions of x , for $\lambda = 0.7$ and $\mu = 0.4$. The middle zero is simple (hence $G_+ \not\equiv 0$) and the corresponding x is not in the spectrum.

However, as we show later there exists just one transcendental function such that x belongs to the spectrum if and only if it is a zero of this single function. This is a desirable property in a general method for finding spectra of systems described in Bargmann representation. An example of the Rabi model with broken $\mathbb{Z}_2$ symmetry, given in the end of paper [1] shows that a direct generalisation of the Braak approach is not obvious.

In the Bargmann representation many models are described by a system of linear differential equations with rational coefficients. The system depends on a spectral parameter E . The problem is to distinguish all values E_n of E such that for $E = E_n$, the considered system possesses a non-zero entire solution contained in the Cartesian product of an appropriate number of Bargmann spaces $\mathcal{H}$.

Here, for simplicity, we consider a system described by just one equation of the second order

$$\varphi'' + p(z)\varphi' + q(z)\varphi = 0, \quad (12)$$

where $p(z)$ and $q(z)$ are rational functions z , which additionally may depend rationally on some parameters. We assume that the system possesses regular singular points $s_1, \dots, s_m \in \mathbb{C}$. Infinity may be a regular or an irregular singular point. Let ρ_i and σ_i be the exponents at singular point s_i . We want to formulate necessary conditions for the existence of a nonzero entire solution of equation (12). Hence, let us assume that $\varphi(z)$ is such a solution. Thus, at each point s_i one of exponents, let us say ρ_i , is a non-negative integer, and a local holomorphic solution at this point has the form

$$\varphi_i(z) := \sum_{n=0}^{\infty} a_n^{(i)} (z - s_i)^{\rho_i + n}. \quad (13)$$

The radius of convergence of this series is not smaller than $r_i := \min_{j \neq i} |s_i - s_j|$. Coefficients $a_n^{(i)}$ can be computed recursively.

To proceed we have to make the following technical assumption. Namely, we require that

$$U_i := D(s_i, r_i) \cap D(s_{i+1}, r_{i+1}) \neq \emptyset, \quad (14)$$

for $i = 1, \dots, m-1$. If none of σ_i is a non-negative integer, then, up to a multiplicative constant, series (13) are just Taylor expansions of entire solution $\varphi(z)$ at points s_i . This is why local solutions $\varphi_i(z)$ and $\varphi_{i+1}(z)$ have to be proportional for all $z \in U_i$ where both are well defined. Hence, for a certain nonzero $\alpha_i \in \mathbb{C}$, the function $F_i(z) := \varphi_i(z) - \alpha_i \varphi_{i+1}(z)$ vanishes for all $z \in U_i$, so it vanishes identically. But $F_i(z)$, as a linear combination of solutions of equation (12), is its solution. So, $F_i(z)$ vanishes identically if and only if $F_i(z_*) = 0$, and $F'_i(z_*) = 0$, for an arbitrary $z_* \in U_i$. Eliminating α_i from equations $F_i(z_*) = 0$ and $F'_i(z_*) = 0$, we obtain the following criterion. If, with the given assumptions, equation (12) has a non-zero entire solution, then

$$W_i(z_i) := \det \begin{bmatrix} \varphi_i(z_i) & \varphi_{i+1}(z_i) \\ \varphi'_i(z_i) & \varphi'_{i+1}(z_i) \end{bmatrix} = 0, \quad (15)$$

for a certain $z_i \in U_i$, and $i = 1, \dots, m-1$. These $(m-1)$ conditions guarantee that $\varphi(z)$ is an entire function. If infinity is a regular singular point, they guarantee also that $\varphi(z)$ belongs to the Bargmann space.

Formulation of this method with relaxed assumptions and its version valid for systems of differential equations will be published in [10].

Let us consider the $\mathbb{Z}_2$ symmetry-broken Rabi model (2). An elimination of $\psi_2(z)$ from this system gives one second order equation for $\varphi(z) := \psi_1(z)$. This equation has the form (12) with

$$p(z) = -\frac{\lambda + 2\epsilon\lambda + z(-1 + 2E + 2\lambda^2)}{z^2 - \lambda^2} \\ q(z) = -\frac{\epsilon^2 - E^2 + 2z\epsilon\lambda + \lambda(\lambda + z(-1 + z\lambda)) + \mu^2}{z^2 - \lambda^2}. \quad (16)$$

Then, we introduce variables y and $v(y)$ setting $z = \lambda(2y - 1)$, and

$$v(y) := \exp(-2\lambda^2 y) \psi_1(\lambda(2y - 1)). \quad (17)$$

By direct computation we obtain that $v(y)$ satisfies the following equation

$$v'' + \left(\alpha + \frac{\beta + 1}{y} + \frac{\gamma + 1}{y - 1} \right) v' + \left(\frac{\tilde{\mu}}{y} + \frac{\tilde{\nu}}{y - 1} \right) v = 0, \quad (18)$$

where

$$\tilde{\mu} = \frac{1}{2}(\alpha - \beta - \gamma + \alpha\beta - \beta\gamma) - \eta, \\ \tilde{\nu} = \frac{1}{2}(\alpha + \beta + \gamma + \alpha\gamma + \beta\gamma) + \delta + \eta,$$

and parameters $a_0 = (\alpha, \beta, \gamma, \delta, \eta)$ were defined earlier, see (6). Equation (18) is the Heun confluent equation. Its local holomorphic solution near $y = 0$ is given by the Heun confluent function $v_0(y) = \text{HeunC}(a_0; y)$. Solution holomorphic near $y = 1$ is given by $v_1(y) = \text{HeunC}(a_1, 1-y)$, where $a_1 := (-\alpha, \gamma, \beta, -\delta, \delta+\eta)$. Hence, local solutions $\varphi_{\mp}(z)$ holomorphic near singular points $z = \mp\lambda$ are given by

$$\varphi_{-}(z) := \exp[\lambda z + \lambda^2] \text{HeunC}(a_0; (1 + z/\lambda)/2) \quad (19)$$

and

$$\varphi_{+}(z) := \exp[\lambda z + \lambda^2] \text{HeunC}(a_1; (1 - z/\lambda)/2). \quad (20)$$

These local solutions are given by expansion of one entire function, and hence their Wronskian

$$W(p) := \varphi_{+}(z_{\star})\varphi'_{-}(z_{\star}) - \varphi'_{+}(z_{\star})\varphi_{-}(z_{\star}), \quad (21)$$

must vanish at arbitrary $z_{\star}$. The graph of the Wronskian $W(p)$ as a function of energy for the Rabi model and the generalised Rabi model are shown in Fig. 3, and Fig. 4, respectively. The resulting spectrum as seen in Fig. 5 reproduces that of [1], with the benefit of not including possible isolated zeroes of G_{+} .




FIG. 3. Graph of Wronskian $W(p)$ for $p := (x, \lambda, \mu, \epsilon) = (x, 7/10, 4/10, 0)$.




FIG. 4. Graph of Wronskian $W(p)$ for $p := (E, \lambda, \mu, \epsilon) = (E, 7/10, 4/10, 2/10)$.

This research has been partially supported by grant No. DEC-2011/02/A/ST1/00208 of National Science Centre of Poland.




FIG. 5. Spectrum of generalised Rabi model for $\mu = 0.7$, and $\epsilon = 0.2$.

* maciejka@astro.ia.uz.zgora.pl

† M.Przybylska@proton.if.uz.zgora.pl

‡ stachowiak@cft.edu.pl

- [1] D. Braak, Phys. Rev. Lett. **107**, 100401 (2011).
- [2] D. Braak, ArXiv e-prints (2012), arXiv:1205.3439 [quant-ph].
- [3] A. Moroz, ArXiv e-prints (2012), arXiv:1205.3139 [quant-ph].
- [4] V. Bargmann, Comm. Pure Appl. Math. **14**, 187 (1961).
- [5] S. Schweber, Annals of Physics **41**, 205 (1967).
- [6] W. Gautschi, SIAM Rev. **9**, 24 (1967).
- [7] A. Ronveaux, *Heun's Differential Equations* (Oxford University Press, Oxford, 1995).
- [8] D. Braak, "Supplement material to paper "integrability of the Rabi model", "<http://prl.aps.org/supplemental/PRL/v107/i10/e100401>" (2011).
- [9] M. Kuś and M. Lewenstein, J. Phys. A **19**, 305 (1986).
- [10] M. Kuś, A. J. Maciejewski, M. Przybylska, and T. Stachowiak, (2012), in preparation.